

The Evening Sky Map

FREE* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

WWW.SKYPMAPS.COM

SOUTHERN HEMISPHERE
FEBRUARY 2007

Sky Calendar – February 2007

- Full Moon** at 5:45 UT. The full Moon of February is called the “Snow Moon”, “Hunger Moon” or “Wolf Moon”.
- Moon very near Saturn** at 24h UT (midnight sky). Occultation visible from Scandinavia, Russia, central Asia, and w. India.
- Moon very near Regulus** at 15h UT (morning sky). Occultation visible from NW North America.
- Mercury near Venus** at 23h UT (evening sky). Look for elusive Mercury about 6° west of Venus above the western horizon. Mags. -3.9 and -0.8.
- Moon at apogee** (farthest from Earth) at 13h UT (distance 404,992 km; angular size 29.5').
- Mercury at greatest elongation**, 18° east from Sun (evening sky) at 18h UT. Mag. -0.5, low in the west (look below Venus after sunset).
- Venus 0.7° SE of Uranus** at 20h UT (evening sky). Mags. -3.9 and +5.9.
- Moon very near Spica** at 2h UT (morning sky).
- Last Quarter Moon** at 9:51 UT.
- Saturn at opposition** (opposite the Sun) at 18h UT. Visible all night long, the ringed planet is at its brightest (mag. +0.0) and closest (disk diameter 20.3") all year. Saturn's rings appear magnificent even in a small telescope.
- Moon very near Antares** at 22h UT (morning sky).
- Moon near Mars** at 3h UT (morning sky). Mag. +1.3.
- New Moon** at 16:14 UT. Beginning of lunation 1041.
- Moon at perigee** (closest to Earth) at 10h UT (361,434 km; 33.1').
- Moon near Venus** at 16h UT (evening sky).
- Moon near the Pleiades** at 23h UT (evening sky).
- First Quarter Moon** at 7:56 UT.
- Moon near Pollux** at 23h UT (evening sky).


Comet McNaught (C/2006 P1) is now a circumpolar object for sky watchers in the southern hemisphere (see sky map). The comet will fade significantly during the month as it moves further from the Sun each day. A spectacular naked-eye object in January, a pair of binoculars or telescope will soon be required to observe the now famous Comet McNaught.

All times in Universal Time (UT). (Australian Eastern Daylight Time = UT + 11 hrs.)



SAVE ON RECOMMENDED PRODUCTS • <http://Skymaps.com/store>

- ★ STAR ATLASES & PLANISPHERES
 - ★ STAR CHARTS & ASTRO POSTERS
 - ★ BOOKS FOR SKY WATCHERS
 - ★ LARGE FORMAT HUBBLE PRINTS
- All sales support the production and free distribution of The Evening Sky Map.



SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY FEB 10 PM

LATE FEB 9 PM

(Add 1 Hour For Daylight Saving)

SKY MAP DRAWN FOR

A LATITUDE OF 35°

SOUTH AND IS

SUITABLE FOR

LATITUDES UP

TO 15° NORTH

OR SOUTH

OF THIS

Symbols

- Galaxy ☾
- Double Star ●●
- Variable Star ⊙
- Diffuse Nebula □
- Planetary Nebula ◇
- Open Star Cluster ○
- Global Star Cluster ⊕

Star Magnitudes ●●●●●
-1 0 1 2 3 4

Copyright © 2000–2007 Kym Thalassoudis. All Rights Reserved.

* TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Australian Eastern Standard Time (for example Sydney, Australia) is 10 hours ahead of UT.

Variable Star – A star that changes brightness over a period of time.

SOUTHERN HEMISPHERE FEBRUARY 2007 CELESTIAL OBJECTS



Easily Seen with the Naked Eye

Sirius	CMa	• The brightest star in the sky. Also known as the "Dog Star". Dist=8.6 ly.
Procyon	CMi	• Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
Canopus	Car	• Second brightest star in the sky. 14,000 times more luminous than the Sun. Dist=310 ly.
β Centauri	Cen	• With Alpha Centauri, forms the so-called "Pointers-to-the-Cross". Dist=525 ly.
α Centauri	Cen	• Nearest bright star to Sun at 4.4 ly. Brilliant double star in a telescope. 80 year period.
Achernar	Eri	• Brightest star in Eridanus, The River. Arabic name meaning "end of river". Dist=144 ly.
Castor	Gem	• Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	• With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
Regulus	Leo	• Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Rigel	Ori	• The brightest star in Orion. Blue supergiant star with mag 7 companion. Dist=770 ly.
Betelgeuse	Ori	• One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Pleiades	Tau	• The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=380 ly.
Hyades	Tau	• Large V-shaped star cluster. Binoculars reveal many more stars. Dist=151 ly.
Aldebaran	Tau	• Brightest star in Taurus. It is not associated with the Hyades star cluster. Dist=65 ly.

Easily Seen with Binoculars

M44	Cnc	• Praesepe or Beehive Cluster. Visible to the naked eye. Dist=577 ly.
M41	CMa	• First recorded observation by Aristotle in 325 BC as "cloudy spot". Dist=2,300 ly.
2516	Car	• Spectacular open star cluster of 100 stars spanning 1/2 deg. Dist=1,300 ly.
2808	Car	• Located 4 deg W of Nu Carinae. Visible to the naked eye on clear nights.
3114	Car	• Stunning open cluster. 30+ stars visible through 7x binoculars. Dist=2,900 ly.
3293	Car	• Rich, tightly packed. Surrounded by large, faint nebulosity. Dist=8,500 ly.
IC 2602	Car	• The "Five of Diamonds". Bright cluster twice diameter of full Moon. Dist=500 ly.
3372	Car	□ Eta Carinae Nebula. Enormous glowing cloud in rich star field. Dist=8,000 ly.
3532	Car	• Herschel - "most brilliant cluster". 60+ stars in 7x binoculars. Dist=1,300 ly.
ω Centauri	Cen	• Largest and brightest globular star cluster in sky. 1 million stars. Dist=17,000 ly.
4755	Cru	• Jewel Box. Outstanding star cluster. Many contrasting colours. Dist=7,600 ly.
LMC	Dor	• Large Magellanic Cloud. A neighbouring galaxy of the Milky Way. Dist=180,000 ly.
M35	Gem	• Fine open cluster located near foot of the twin Castor. Dist=2,800 ly.
γ Leporis	Lep	• Visible with binoculars. Gold & white stars. Mags 3.6 & 6.2. Dist=30 ly. Sep=96.3".
2232	Mon	• A large scattered star cluster of 20 stars. Dist=1,300 ly.
2244	Mon	• Surrounded by the rather faint Rosette Nebula. Dist=5,540 ly.
M50	Mon	• Visible with binoculars. Telescope reveals individual stars. Dist=3,000 ly.
M42	Ori	□ The Great Orion Nebula. Spectacular bright nebula. Best with telescope. Dist=1,500 light years.
L ²	Pup	• Semi-regular variable. Magnitude varies between 2.6 & 6.2 over 140.42 days.
M47	Pup	• Bright star cluster. 15+ stars in 7x binoculars. Dist=1,500 ly.
M46	Pup	• Dist=5,400 ly. Contains planetary NGC 2438 (Mag 11, d=65") - not associated.
2451	Pup	• 30+ stars in binoculars. The brightest star, c Puppis, is red. Dist=850 ly.
2477	Pup	• Very rich but distant star cluster (4,200 ly). Resembles globular through binoculars.
47 Tucanae	Tuc	• Spectacular object. Telescope will reveal stars. Near edge of SMC. Dist=15,000 ly.
β Tucanae	Tuc	• Complex multiple star. Binoculars show one pair. Telescope required to split primary star.
SMC	Tuc	• Small Magellanic Cloud. Companion galaxy to Milky Way. Requires dark sky. Dist=210,000 ly.
2547	Vel	• Fine open cluster visible through binoculars. Dist=1,300 ly.
IC 2391	Vel	• Omicron Velorum Cluster. Superb object for binoculars. Dist=450 ly.

Telescopic Objects

M67	Cnc	• Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.
3918	Cen	• The Blue Planetary. Visible in a small telescope as a round blue disk.
2070	Dor	□ Tarantula Nebula. A bright nebula located in LMC. A star-forming region.
θ Eridani	Eri	• Striking blue-white double star. Mags 3.2 & 4.3. Visible in a small telescope. Sep=8.2".
3242	Hya	• Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.
β Monocerotis	Mon	• Triple star. Mags 4.6, 5.0 & 5.4. Requires telescope to view arc-shape. Sep=7.3".
2264	Mon	• Christmas Tree Cluster. Associated with the Cone Nebula. Dist=2,450 ly.
σ Orionis	Ori	• Superb multiple star. 2 mag 7 stars one side, mag 9 star on other. Struve 761 triple in field.
k Puppis	Pup	• Telescope easily shows two blue-white stars of almost equal brightness. Sep=9.9".
M1	Tau	□ Crab Nebula. Remnant from supernova which was visible in 1054. Dist=6,500 ly.
3132	Vel	• One of the brightest planetaries. Magnitude 10 central star. Dist=2,600 ly.